



Above is a photo of Admelog (insulin lispro injection) pen.

What is Admelog (insulin lispro injection)?

Prescription Admelog is a fast-acting human insulin used to improve blood sugar control in adults with Type 2 diabetes and adults and children (3 years and older) with Type 1 diabetes.

Important Safety Information for Admelog (insulin lispro injection)

Do not use Admelog if you have low blood sugar or if you are allergic to insulin lispro or any of the other ingredients in Admelog.

Do not share needles, insulin pens, or syringes with others, even if the needle is changed. Do NOT reuse needles. You may give other people a serious infection, or get a serious infection from them.

Before starting Admelog, tell your doctor about all your medical conditions, including if you have liver or kidney problems, if you are pregnant or planning to become pregnant or if you are breastfeeding or planning to breastfeed.

Heart failure can occur if you are taking insulin together with pills called TZDs (thiazolidinediones), even if you have never had heart failure or other heart problems. If you have heart failure, it may get worse while you take TZDs with Admelog. Your treatment with TZDs and Admelog may need to be changed or stopped by your doctor if you have new or worsening heart failure. Tell your doctor if you have any new or worsening symptoms, including:

- Shortness of breath
- Sudden weight gain
- Swelling of your ankles or feet

Tell your doctor about all the medications you take, including OTC medicines, vitamins, supplements, and herbal supplements.

When used as a mealtime insulin, Admelog should be taken within 15 minutes before a meal or immediately after a meal. Test your blood sugar levels as directed by your doctor when using Admelog. Do not change your dose or type of insulin without talking to your doctor. Any change of insulin should be made cautiously and only under medical supervision.

Check your insulin label to verify that you have the correct insulin before each injection. Admelog must only be used if the solution is clear and colorless with no particles visible.

Please see additional Important Safety Information for Admelog on the following page.

Please see full [Prescribing Information](#).

IMPORTANT SAFETY INFORMATION FOR ADMELOG (insulin lispro injection) (continued)

While using Admelog, do not drive or operate heavy machinery until you know how Admelog affects you. Do not drink alcohol or use medicines that contain alcohol.

Admelog may cause serious side effects that can lead to death, including:

- **Low blood sugar (hypoglycemia).** Signs and symptoms of low blood sugar may include dizziness or lightheadedness, sweating, confusion, headache, blurred vision, slurred speech, shakiness, fast heartbeat, anxiety, irritability or mood changes, hunger.
- **Serious allergic reactions.** Get medical help right away if you have: a rash over your whole body, trouble breathing, fast heartbeat, feeling faint, or sweating.
- **Low potassium in your blood (hypokalemia).** Your doctor should monitor you for low blood potassium levels.
- **Heart failure.** Your doctor should monitor you closely while you are taking TZDs with Admelog.

Get emergency medical help if you have:

- Trouble breathing, shortness of breath, fast heartbeat, swelling of your face, tongue, or throat, sweating, extreme drowsiness, dizziness, confusion.

Admelog may have additional side effects including injection site reactions which may include change in fat tissue, skin thickening, redness, swelling and itching.

Important Safety Information for Admelog (insulin lispro injection) SoloStar

Admelog SoloStar is a disposable prefilled insulin pen. Talk to your doctor about proper injection technique and follow instructions in the *Instruction Leaflet* that comes with the pen.

Please see full [Prescribing Information](#).

SAUS.ILI.17.11.9057(1)