



Above is a photo of a Toujeo® (insulin glargine injection) 300 Units/mL SoloSTAR® pen.

What is Toujeo® (insulin glargine injection) 300 Units/mL?

Prescription Toujeo® is a long-acting insulin used to control blood sugar in adults with diabetes mellitus.

- Toujeo® contains 3 times as much insulin in 1 mL as standard insulin (100 Units/mL)
- Toujeo® is not for use to treat diabetic ketoacidosis
- Toujeo® should not be used in children

Important Safety Information for Toujeo® (insulin glargine injection) 300 Units/mL

Do not take Toujeo® if you have low blood sugar or if you are allergic to insulin or any of the ingredients in Toujeo®.

Do NOT reuse needles or share insulin pens even if the needle has been changed. Do not share your Toujeo® SoloStar® or Toujeo® Max SoloStar® pen with other people, even if the needle has been changed. You may give other people a serious infection, or get a serious infection from them. Reusing needles increases your chance of having blocked needles.

Before starting Toujeo®, tell your doctor about all your medical conditions, including if you have liver or kidney problems, if you are pregnant or planning to become pregnant or if you are breastfeeding or planning to breastfeed.

Heart failure can occur if you are taking insulin together with pills called TZDs (thiazolidinediones), even if you have never had heart failure or other heart problems. If you have heart failure, it may get worse while you take TZDs with Toujeo®. Your treatment with TZDs and Toujeo® may need to be changed or stopped by your doctor if you have new or worsening heart failure. Tell your doctor if you have any new or worsening symptoms including:

- Shortness of breath
- Sudden weight gain
- Swelling of your ankles or feet

Tell your doctor about all the medications you take, including OTC medicines, vitamins, and supplements, and herbal supplements.

Toujeo® should be taken at the same time once a day. Test your blood sugar levels daily while using any insulin, including Toujeo®. Do not change your dose or type of insulin without talking to your doctor. Verify you have the correct insulin before each injection.

Please see additional Important Safety Information for Toujeo® on the following page.

Please see full [Prescribing Information](#).

**IMPORTANT SAFETY INFORMATION FOR TOUJEO® (insulin glargine injection)
(continued)**

Do NOT use a syringe to remove Toujeo® from your SoloStar® pen or Toujeo® Max SoloStar® pen. This can cause you to give yourself too much insulin. Toujeo® has 3 times as much insulin in 1 mL compared to other standard insulin pens. Your dose for Toujeo®

may be different from other insulins you have taken. Any change of insulin should be made cautiously and only under medical supervision.

Do NOT dilute or mix Toujeo® with any other insulin or solution. It will not work as intended and you may lose blood sugar control, which could be serious. Use Toujeo® only if the solution is clear and colorless with no particles visible.

While using Toujeo®, do not drive or operate heavy machinery until you know how Toujeo® affects you. Don't drink alcohol or use other medicines that contain alcohol.

The most common side effect of any insulin, including Toujeo®, is low blood sugar (hypoglycemia), which may be serious and can be life-threatening. Severe hypoglycemia may cause harm to your heart or brain. Symptoms of serious low blood sugar may include shaking, sweating, fast heartbeat, and blurred vision.

Toujeo® may cause severe allergic reactions that can lead to death. Get medical help right away if you have:

- A rash over your whole body
- Shortness of breath
- Swelling of your face, tongue, or throat
- Extreme drowsiness, dizziness, or confusion
- Trouble breathing
- Fast heartbeat
- Sweating

Toujeo® may have additional side effects including swelling, weight gain, low potassium, and injection site reactions which may include change in fat tissue, skin thickening, redness, swelling, and itching.

Toujeo® SoloStar® and Toujeo® Max SoloStar® are disposable prefilled insulin pens. It is important to perform a safety test when using a new pen for the first time. Talk to your doctor about proper injection technique and follow instructions in the *Instruction Leaflet* that comes with the pens.

Please see full [Prescribing Information](#).

SAUS.TJO.17.05.2906(1)